

**DRUG FACTS**

Active ingredient(s)	Purpose
.....	.....
.....	.....
.....	.....
.....	.....
.....	.....
.....	.....
.....	.....
.....	.....

**Use(s)**

**Warnings**

Warnings:  
For external use only  
Warning: For external use only.  
Warnings:  
For external use only.  
Warning:  
If you experience eye pain, changes in vision, continued redness or irritation of the eye, or if the condition worsens or persists for more than 72 hours, discontinue use and consult a doctor.  
Do not use if solution changes color or becomes cloudy.  
Warnings: For external use only.  
Avoid contact with eyes. If this happens, rinse thoroughly with water.  
Warnings  
Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include: shock, facial swelling, asthma (wheezing) rash, skin reddening, blisters, hives If an allergic reaction occurs, stop use and seek medical help right away.  
Stomach bleeding warning: This product contains a nonsteroidal anti-inflammatory drug (NSAID), which may cause severe stomach bleeding. The chance is higher if you: are age 60 or older, have had stomach ulcers or bleeding problems, take a blood thinner (anticoagulant) or steroid drug, take other drugs containing NSAIDs (aspirin, ibuprofen, naproxen, or others), have 3 or more alcoholic drinks every day while using this product, take more or for a longer time than directed  
Warnings:  
Liver warning: This product contains acetaminophen.  
Severe liver damage may occur if: adult takes more than 12 tablets in 24 hours, which is the maximum daily amount; child takes more than 5 doses in 24 hours; taken with other drugs containing acetaminophen; adult has 3 or more alcoholic drinks every day while using this product  
Warnings  
Reye's syndrome: Children and teenagers who have or are recovering from chicken pox of flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.  
Allergy alert: Aspirin may cause a severe allergic reaction which may include: hives, skin reddening, facial swelling, rash, asthma (wheezing), blisters, shock, If an allergic reaction occurs, stop use and seek medical help right away.  
Stomach bleeding warning: This contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you: are age 60 or older; have had stomach ulcers or bleeding problems; take a blood thinner (anticoagulant) or steroid drug; take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others); have 3 or more alcoholic drinks every day while using this product; take more or for a longer time than directed  
Warnings: For external use only.  
Warnings:  
For external use only  
Flammable, keep away from fire or flame

**Do not use**

Do not use: In eyes, in large quantities, over raw blistered areas, or on deep puncture wounds, animal bites or serious burns, for more than one week  
Do not use in the eyes or over large areas of the body.  
Do not use: in eyes; over large areas of the body;  
If allergic to any of the ingredients; for more than one week unless directed by a physician.  
Do not use: In eyes, on broken skin, deep puncture wounds. If unusual redness, swelling, irritation or other symptoms occur, consult a physician immediately.  
Do not use if you have ever had an allergic reaction to any other pain reliever/fever reducer, right before or after heart surgery.  
Do not use: with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist; for more than 10 days for pain unless directed by a doctor; for more than 3 days for fever unless directed by a doctor  
Do not use: if you have ever had an allergic reaction to any other pain reliever/ fever reducer; right before or after heart surgery; if you are taking a prescription drug for gout, diabetes or arthritis  
Do not use: with electrocautery, in the eyes.  
Do not use in the eyes. In case of contact, rinse eyes thoroughly with water.

**Ask a doctor before use if you have**

Ask a doctor before use if stomach bleeding warning applies to you; you have a history of stomach problems such as heartburn; you have a high blood pressure, heart disease, liver cirrhosis, or kidney disease; you are taking a diuretic  
Ask a doctor before use if the user has liver disease  
Ask a doctor before use if: stomach bleeding warning applies to you; you have a history of stomach problems such as heartburn; you have high blood pressure, heart disease, liver cirrhosis, or kidney disease; you are taking a diuretic

**Ask a doctor or pharmacist before use if you are**

Ask a doctor before use if you are:  
taking any other drug containing NSAID (prescription or nonprescription); taking aspirin for heart attack or stroke, because Ibuprofen may decrease this benefit of aspirin; taking any other drug  
Ask a doctor or pharmacist before use if the user is taking the blood thinning drug warfarin  
Ask a doctor or pharmacist before use if you are:  
under a doctor's care for any serious condition; taking any other drug

**When using this product**

Directions: Tear open packet, unfold towelette and use to cleanse desired skin area. Discard towelette appropriately after single use.  
Directions: clean affected area; apply small amount of this product (an amount equal to the surface area of the tip of a finger) on the area 1 to 3 times daily; may be covered with a sterile bandage  
When using this product:  
take with food or milk if stomach upset occurs  
When using this product: take with food or milk if stomach upset occurs  
Directions  
do not use more than directed  
the smallest effective dose should be used  
drink a full glass of water with each dose  
do not take longer than 10 days, unless directed by a doctor  
Wipe Injection site vigorously and discard.

**Stop use and ask a doctor if**

Stop use if unusual redness, swelling or other symptoms occur. Consult a physician immediately.  
Stop use and consult a doctor:  
if the condition persists or gets worse; a rash or other allergic reaction develops  
Stop use and ask a doctor If:  
you experience any of the following signs of stomach bleeding; feel faint; vomit blood; have bloody or black stools; have stomach pain that does get better; pain gets worse or lasts more than 10 days; fever gets worse or lasts more than 3 days; redness or swelling is present in the painful area; any new symptoms appear

Stop use and ask a doctor if: symptoms do not improve; pain gets worse or lasts for more than 10 days; fever gets worse or lasts for more than 3 days; new symptoms occur; redness or swelling is present; a rare sensitivity reaction occurs

Stop use and ask a doctor if you experience any of the following signs of stomach bleeding:

feel faint; vomit blood; have bloody or black stools; have stomach pain that does not get better; pain gets worse or lasts more than 10 days; fever gets worse or lasts more than 3 days; you have difficulty swallowing; if ringing in the ears or loss of hearing occurs; redness or swelling is present in the painful areas; any new symptoms appear

Stop use if irritation and redness develop. If condition persists for more than 72 hours, consult your doctor.

Stop use and consult a doctor if irritation and redness develop and persist for more than 72 hours.

**If pregnant or breast-feeding**

**Keep out of reach of children**

Keep out of reach of Children.

If ingested, contact a Poison Control Center right away.

Keep out of reach of children: If swallowed, get medical help or contact a Poison Control Center right away.

Keep out of reach of children.

If ingested, contact a Poison Control Center right away.

Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Keep out of reach of children. In case of accidental overdose, contact a doctor or Poison Control Center immediately. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms. Do not exceed recommended dosage

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.

Keep out of reach of children.

If swallowed, get medical help or contact a Poison Control Center right away.

***Directions***

Directions: Clean affected area, Apply small amount not more than 3 times daily.

May be covered with a sterile bandage.

Directions

Remove contacts before using.

Twist top to remove.

Flush the affected area as needed. Control

Rate of flow by pressure on the bottle. Do not touch tip of the container to any surface. Do not reuse.

If necessary continue flushing with emergency eyewash or shower.

Discard bottle after use.

Directions:

do not use more than directed; the smallest effective dose should be used; do not take longer than 10 days, unless directed by a doctor.

Adults and Children (12 years and older): Take 1 tablet every 4 to 6 hours while symptoms persist. If pain or fever does not respond to 1 tablet, 2 tablets may be used. Do not exceed 6 tablets in 24 hours, unless directed by a doctor.

Children under 12 years: Do not give to children under 12 years of age.

Directions

Adults and Children 12 years of age or older	Take 2 tablets every 4 to 6 hours as needed. Do not take more than 12 tablets in 24 hours.
Children 6-11 years of age	Take 1 tablet every 4 to 6 hours as needed. Do not take more than 5 tablets in 24 hours.
Children under 6 years of age	Do not use this regular strength product. This will provide more than the recommended dose (overdose) and could cause serious health problems.

Adults and children: (12 years and older) Take 1 or 2 tablets with water every 4 hours as needed. Do not take more than 12 tablets in 24 hours, or as directed by a doctor.

Children under 12 years: Do not give to children under 12 years of age.

Directions:

Empty contents into palm. Rub hands until gel dissipates. Recommended for repeated use.

***Other information***

Other Information:

Store at room temperature

Other information:

Store at room temperature.

Store in a cool place. For irrigation only.

Discard unused portion of the solution.

Not for injection.

Flammable - keep away from fire or flame.

Other information: Store at controlled room temperature; avoid excessive heat 40 degree Celsius (104 degree Fahrenheit); tamper evident sealed packets; do not use any opened or torn packets

Store at 59-86 degree Fahrenheit (15-30 degree Celsius)

Store at 59 - 86 degree Fahrenheit (15 - 30 degree Celsius); avoid excessive heat and humidity; tamper evident sealed packets;

Do not use any opened or torn packets

Flammable - keep away from fire or flame

Store at room temperature 15-30 degree Celsius (59-86 degree Fahrenheit)

Other Information: Store at 15 to 25 degree Celsius (59 to 77 degree Fahrenheit)

***Inactive ingredients***

Inactive Ingredient: Purified water

Inactive Ingredients:

Sodium Chloride USP	44mg
Monobasic Sodium Phosphate USP	18mg
Sodium Phosphate Dibasic USP	111mg
Edetate Disodium USP	10mg
Benzalkonium Chloride	0.5mg
NF (as preservative)	

Inactive ingredients: cellulose, corn starch, fumed silica gel, hypromellose, lactose, magnesium stearate, polydextrose, polyethylene glycol, povidone, silica, sodium starch glycolate, stearic acid, titanium dioxide, triacetin.

Inactive Ingredients: hypromellose, polyethylene glycol, propylene glycol, corn starch  
Inactive Ingredient: Purified water.